

All Algebraic Functions Can Be Computed Fast

H. T. KUNG AND J. F. TRAUB

Carnegie-Mellon University, Pittsburgh, Pennsylvania

ABSTRACT The expansions of algebraic functions can be computed "fast" using the Newton Polygon Process and any "normal" iteration. Let $M(j)$ be the number of operations sufficient to multiply two j th-degree polynomials. It is shown that the first N terms of an expansion of any algebraic function defined by an n th-degree polynomial can be computed in $O(nM(N))$ operations, while the classical method needs $O(N^n)$ operations. Among the numerous applications of algebraic functions are symbolic mathematics and combinatorial analysis. Reversion, reciprocation, and n th root of a polynomial are all special cases of algebraic functions.

KEY WORDS AND PHRASES computational complexity, fast computation, symbolic mathematics, combinatorial mathematics, Newton Polygon, Newton iteration, algebraic functions, reversion

CR CATEGORIES 5.15, 5.25, 5.7

1 Introduction

Let

$$P(W, z) = A_n(z)W^n + \cdots + A_0(z), \quad (1.1)$$

where the $A_i(z)$ are polynomials over a field Δ . In general we shall take Δ to be the field of complex numbers; Section 7 is an exception. (Many of the results hold for an algebraically closed field of characteristic 0.) Without loss of generality we assume $A_0(z) \neq 0$ and $A_n(z) \neq 0$. Usually, but not always, capital letters will denote polynomials or series; lower-case letters will denote scalars.

The zero of (1.1), a function $S(z)$ such that $P(S(z), z) \equiv 0$, is called the *algebraic function* corresponding to $P(W, z)$. Let z_0 be an arbitrary complex number, finite or infinite. It is known from the general theory of algebraic functions that $S(z)$ has n fractional power series expansions around z_0 . By the *computation of an algebraic function* we shall mean the computation of the first N coefficients (including zero coefficients) of one of its expansions. (This will be made precise in Section 3.) The problem we study in this paper is the computation of one expansion of the algebraic function. Our results can be modified for computing more than one expansion or all expansions of the algebraic function.

As described in most texts, the classical method computes algebraic functions by comparison of coefficients. It is not difficult to show that the method can take $O(N^n)$ operations, where n is the degree of $P(W, z)$ with respect to W . Hence the classical method is very slow when n is large.

General permission to make fair use in teaching or research of all or part of this material is granted to individual readers and to nonprofit libraries acting for them provided that ACM's copyright notice is given and that reference is made to the publication, to its date of issue, and to the fact that reprinting privileges were granted by permission of the Association for Computing Machinery. To otherwise reprint a figure, table, other substantial excerpt, or the entire work requires specific permission as does republication, or systematic or multiple reproduction.

The major results of this paper were announced at the Symposium on Algorithms and Complexity: New Directions and Recent Results, Carnegie-Mellon University, April 7-9, 1976 (abstract in [11]).

This work was supported in part by the National Science Foundation under Grant MCS 75-222-55 and the Office of Naval Research under Contract N00014-76-C-0370, NR 044-422.

Authors' address: Department of Computer Science, Carnegie-Mellon University, Pittsburgh, PA 15213

© 1978 ACM 0004-5411/78/0400-0245 \$00.75

The main result of this paper is that *every algebraic function can be computed fast*. Let $M(N)$ denote the number of operations sufficient to multiply two N th-degree polynomials over the field Δ . Let $C(N)$ be the number of operations needed to compute any algebraic function. We prove that $C(N) = O(nM(N))$. Since $M(N) = O(N^2)$ (or $M(N) = O(N \log N)$ if the FFT is used), our algorithms are considerably faster than the classical method even for moderate n . It is an open problem whether or not a general algebraic function can be computed in less than $O(M(N))$ operations.

The "fast computation" of the title is because the coefficients of a "regular" problem can always be computed fast by iteration (Section 5) and the general problem can be reduced to a regular problem (Section 6) with cost independent of N .

Brent and Kung [5] showed that the cost for reversion of a polynomial, which is a very special case of an algebraic function (see discussion later in this section), is $O((N \log N)^{1/2}M(N))$. We stated above that the cost of expanding an algebraic function is $O(nM(N))$. These results are reconciled by the observation that we are considering the case that the degree n of $P(W, z)$ with respect to W is fixed and independent of N , while Brent and Kung considered the case where $n = N$.

There are known examples of computation using Newton-like iteration in settings such as algebraic number theory [8, 2], power series computation [10, 5], and the Zassenhaus construction in p -adic analysis [21]. Computation of algebraic functions raises certain issues not present in these other settings; see especially Section 6. As we will see in Section 5, there is nothing special about Newton-like iteration; any "normal iteration" can be used.

Although the complexity results are stated asymptotically, Theorems 5.1 and 6.1 give nonasymptotic analyses of the algorithms. *Hence various nonasymptotic analyses can also be carried out.*

We are interested in the computation of algebraic functions for a number of reasons. These include

1. A number of problems where fast algorithms are known are special cases of algebraic functions. (More details are given below.)

2. There are numerous applications. For example, many generating functions of combinatorial analysis and functions arising in mathematical physics are algebraic functions. The integrands of elliptic and more generally Abelian integrals are algebraic functions. See Section 9 for an example.

3. Algorithms for expanding algebraic functions are needed in systems for symbolic mathematics such as MACSYMA [14].

4. Algebraic functions are of theoretical interest in many areas of mathematics. These include integration in finite terms [17], theory of plane curves [20], elliptic function theory [6], complex analysis [1, 18], and algebraic geometry [12]. Algebraic function theory is a major subject in its own right. (See, for example, [3, 7].)

We exhibit special cases of algebraic functions where fast algorithms are known.

A. *Reciprocal of a polynomial.* $P(W, z) = A_1(z)W - 1$. (See Kung [10].) (Actually Kung uses $P(W, z) = W^{-1} - A_1(z)$ which is not of the form (1.1), and allows $A_1(z)$ to be a power series.)

B. *n -th root of a polynomial.* $P(W, z) = W^n - A_0(z)$. (See [4, Sec 13], where the $A_0(z)$ is allowed to be a power series.)

C. *Reversion of a polynomial* Let F be a given polynomial with zero constant term. We seek a function G such that $F(G(z)) = z$. To see that this is a special case of an algebraic function, let $F(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x$. Then we seek $G(x)$ such that $a_n G^n(z) + \dots + a_1 G(z) - z = 0$. This is an instance of our general problem with $A_i(z) = a_i$, $i = 1, \dots, n$, $A_0(z) = -z$. See [5].

We summarize the results of this paper. In Section 2 we show that without loss of generality we can take $z_0 = 0$ and assume $A_n(0) \neq 0$. Notation is established and a few basic facts from algebraic function theory are summarized in Section 3. The concept of normal iteration is introduced in Section 4 and convergence of normal iterations for

regular problems is established in Section 5. In Section 6 we state and analyze the Newton Polygon Process, which reduces the general problem to a regular problem. A symbolic mode of computation with exact arithmetic is introduced in Section 7. Section 8 shows that $C(N) = O(nM(N))$. In Section 9 we give a number of examples, several of which are more general than the theory of the preceding sections. Section 10 discusses extensions of the work presented here.

In this paper we analyze algorithms under the assumption that the coefficient of power series are “nongrowing,” e.g. all coefficient computations are done in a finite field or in finite-precision floating point arithmetic. An analysis dealing with variable-precision coefficients is yet to be performed.

2. Preliminary Transformations

Recall that we wish to compute one of the expansions around z_0 of the algebraic function $S(z)$ corresponding to

$$P(W, z) = A_n(z)W^n + \dots + A_0(z),$$

i.e. $P(S(z), z) \equiv 0$. In this section we show that after two simple transformations we need only deal with the case that $z_0 = 0$ and $A_n(0) \neq 0$. If we transform $P(W, z)$ to $\tilde{P}(W, z)$, then $\tilde{S}(z)$ is defined by $\tilde{P}(\tilde{S}(z), z) \equiv 0$.

Consider first the case $z_0 = \infty$. Let

$$\tilde{P}(W, z) = z^m P(W, 1/z), \tag{2.1}$$

where $m = \max_{0 \leq i \leq n} (\deg A_i)$. By definition, an expansion of $S(z)$ around $z_0 = \infty$ is an expansion of $\tilde{S}(z)$ around $z_0 = 0$.

Consider next the case that z_0 is any finite complex number. Define $\tilde{P}(W, z) = P(W, z + z_0)$. An expansion of $\tilde{S}(z)$ around the origin is an expansion of $S(z)$ around $z = z_0$.

For the remainder of this paper we shall therefore take $z_0 = 0$.

Let $A_n(0) = 0$. Then the algebraic function $S(z)$ corresponding to $P(W, z)$ has one or more expansions with negative powers. Using the following transformation, we need only deal with expansions with nonnegative powers. It is convenient to use ord notation.

Definition 2.1. Let $A(z)$ be an integral or fractional power series. If $A(z) \neq 0$, then $\text{ord}(A)$ denotes the degree of the lowest degree term in $A(z)$. If $A(z) \equiv 0$, then $\text{ord}(A) = \infty$. \square

Choose nonnegative integers μ and λ to satisfy the following conditions:

$$\mu + \text{ord}(A_n) = n\lambda, \quad \mu + \text{ord}(A_i) \geq i\lambda, \quad i = 1, \dots, n - 1.$$

Let $\tilde{P}(W, z) = z^\mu P(W/z^\lambda, z)$. Then the coefficients of $\tilde{P}(W, z)$, $\tilde{A}_i(z)$, are polynomials with $\tilde{A}_n(0) \neq 0$, and $\tilde{S}(z)$ has only expansions with nonnegative powers. Since the expansions of $S(z)$ are those of $\tilde{S}(z)$ divided by z^λ , it suffices to compute expansions of $\tilde{S}(z)$. For the remainder of this paper, we therefore assume that $A_n(0) \neq 0$. (One should note, however, that the results of Section 5 hold without the assumption.)

3. Facts from Algebraic Function Theory

We introduce some notation and state a basic result of algebraic function theory which characterizes the expansions of the algebraic function corresponding to

$$P(W, z) = A_n(z)W^n + \dots + A_0(z).$$

There exist r positive integers d_1, \dots, d_r such that $d_1 + \dots + d_r = n$ and the expansions of the algebraic function are given by

$$S_{i,j}(z) = \sum_{l=0}^{\infty} s_{i,l} \xi_i^{lj} \cdot z^{l/d_i} \tag{3.1}$$

for $i = 1, \dots, r$ and $j = 0, \dots, d_i - 1$, where ξ_i is a primitive d_i th root of unity and the

$s_{i,j}$ are complex numbers. For each i , the expansions $S_{i,j}$, $j = 0, \dots, d_i - 1$, are said to constitute a *cycle*.

The problem considered in this paper is to compute one expansion of an algebraic function. For notational convenience, let the expansion be denoted by

$$S(z) = \sum_{l=0}^{\infty} s_l \cdot z^{l/d}.$$

Hence our problem can be formulated as that of computing the value of d and the coefficients s_0, s_1, \dots . (In this paper $S(z)$ represents either an algebraic function or one of its expansions, depending upon the context.) Note that since $P(S(z), z) \equiv 0$, we have $P(s_0, 0) = 0$. Thus, s_0 is a zero of the numerical polynomial $P(W, 0)$. We say our problem is *regular* with respect to s_0 if s_0 is a simple zero of $P(W, 0)$. (In this definition we allow $A_n(0)$ to be 0.) For a regular problem, we have $d = 1$, that is, the expansion $S(z)$ is an integral power series. In Section 5 we shall show that a regular problem can always be solved by iteration. In Section 6 we shall show how the general problem can be transformed to a regular problem.

4. Normal Iterations

We introduce the concept of a *normal* numerical iteration. We give a novel definition of the order of a normal iteration which is convenient for the application to power series iteration. In Section 5 we shall show that a normal iteration with order greater than unity will always converge if used for a regular problem.

Let $p(w)$ be the numerical polynomial $P(W, 0)$, let s be a zero of $p(w)$, and let $e^{(i)} = w^{(i)} - s$ denote the error of the i th iterate. To motivate the definition of normal iteration we first consider two examples.

Example 4.1 Newton iteration.

$$w^{(i+1)} = w^{(i)} - p(w^{(i)})/p'(w^{(i)}).$$

From the Taylor series expansions

$$p(w^{(i)}) = p'(s)e^{(i)} + (p''(s)/2)(e^{(i)})^2 + \dots$$

and

$$p'(w^{(i)}) = p'(s) + p''(s)e^{(i)} + \dots,$$

we have

$$e^{(i+1)} = (p''(s)/2p'(s))(e^{(i)})^2 + \sum_{j=3}^{\infty} c_j \cdot (e^{(i)})^j, \tag{4.1}$$

where the c_j are rational expressions of the derivatives of p at s , with powers of $p'(s)$ as the denominators. □

Example 4.2. Secant iteration.

$$w^{(i+1)} = w^{(i)} - ((w^{(i)} - w^{(i-1)})/(p(w^{(i)}) - p(w^{(i-1)}))) \cdot p(w^{(i)}).$$

Using the Taylor series expansions of $p(w^{(i)})$ and $p(w^{(i-1)})$, we obtain

$$e^{(i+1)} = (p''(s)/2p'(s))e^{(i)}e^{(i-1)} + \sum_{\substack{j+l \geq 3 \\ j,l \geq 1}} c_{j,l} \cdot (e^{(i)})^j (e^{(i-1)})^l, \tag{4.2}$$

where the $c_{j,l}$ are rational expressions of the derivatives of p at s , with powers of $p'(s)$ as the denominators. □

Consider now a general iteration

$$w^{(i+1)} = \psi(w^{(i)}, w^{(i-1)}, \dots, w^{(i-m)}), \tag{4.3}$$

which is defined in terms of rational expressions of p and its derivatives.

Assume that by using Taylor series expansions, we can derive

$$e^{(t+1)} = \sum_{j=0}^m c_{j_0, \dots, j_m} \cdot (e^{(t)})^{j_0} \dots (e^{(t-m)})^{j_m}, \tag{4.4}$$

where the c_{j_0, \dots, j_m} are rational expressions of the derivatives of p at s .

Definition 4.1. ψ is said to be a *normal iteration* if the denominator of each c_{j_0, \dots, j_m} is a power of $p'(s)$. \square

From (4.1) and (4.2) we have that both Newton iteration and secant iteration are normal. In fact most commonly used iterations are normal. We prove that the classical one-point inverse interpolatory iterations ψ_ρ (see [19, Sec. 5.1]; in particular ψ_2 is the Newton iteration) are normal. Let q denote the inverse function to p and $v^{(t)} = p(w^{(t)})$. Then

$$s = q(0) = q(v^{(t)}) - q'(v^{(t)})v^{(t)} + 1/2 q''(v^{(t)})(v^{(t)})^2 + \dots$$

By definition of ψ_ρ ,

$$s = \psi_\rho(w^{(t)}) + \sum_{j \geq \rho} ((-1)^j/j!)q^{(j)}(v^{(t)})(p(w^{(t)}))^j$$

and

$$e^{(t+1)} = \sum_{j \geq \rho} ((-1)^{j+1}/j!)q^{(j)}(v^{(t)})(p(w^{(t)}))^j.$$

Note that

$$p(w^{(t)}) = p'(s)e^{(t)} + 1/2 p''(s)(e^{(t)})^2 + \dots$$

and that $q^{(j)}(v^{(t)})$ is a rational expression of $p^{(k)}(w^{(t)})$ for $k = 1, \dots, j$ and has the denominator $(p'(w^{(t)}))^j$. Expanding the $p^{(k)}(w^{(t)})$ around s shows that ψ_ρ is a normal iteration.

Definition 4.2. For a normal iteration ψ defined by (4.3) and satisfying (4.4), we define the *order* ρ of ψ by

$$\rho = \sup\{r | r^{m+1} \leq j_0 r^m + j_1 r^{m-1} + \dots + j_m \text{ for all } (j_0, \dots, j_m) \text{ such that } c_{j_0, \dots, j_m} \neq 0 \text{ for some polynomial } p\}. \quad \square$$

By (4.1), it is easy to check that the Newton iteration has order 2. In general it can be shown that the one-point inverse interpolatory iteration ψ_ρ has order ρ . Consider now the secant iteration. By (4.2), the order of the iteration is given by

$$\rho = \sup\{r | r^2 \leq jr + l \text{ for all } j, l \geq 1\},$$

which is equivalent to $\rho = \sup\{r | r^2 \leq r + 1\}$. Hence ρ is the positive root of $r^2 = r + 1$, i.e. $\rho = \phi = (1 + \sqrt{5})/2$.

5 Regular Problems: Normal Iterations on Power Series

We show how normal numerical iterations with order greater than unity can always compute an expansion of an algebraic function for a regular problem. The main result is Theorem 5.1. As a corollary of this theorem we show that a Newton-like iteration always “converges quadratically.” We also show the convergence of a secant-like iteration. We end the section with an example of a convergent *first-order* iteration.

We begin with some definitions. Recall that a meromorphic series is a power series with a finite number of negative powers.

Definition 5.1. Given a meromorphic series $A(z)$ and a real number σ , then by the notation $B(z) \equiv A(z) \pmod{z^\sigma}$, we mean $B(z)$ is a finite series consisting of all terms of $A(z)$ of degree less than σ . \square

Let ψ be a normal numerical iteration. Let the numbers $w^{(t)}, \dots, w^{(t-m)}$ in (4.3), the defining relation for ψ , be replaced by meromorphic series $W^{(t)}(z), \dots, W^{(t-m)}(z)$. Then the iterate $W^{(t+1)}(z)$ defined by

$$W^{(i+1)}(z) = \psi(W^{(i)}(z), \dots, W^{(i-m)}(z))$$

is in general a meromorphic series, provided that it is well defined. Let $E^{(i)}(z) = W^{(i)}(z) - S(z)$ denote the error of the i th iterate.

Definition 5.2. We say an iteration on a meromorphic series *converges* if $\lim_{i \rightarrow \infty} \text{ord}(E^{(i)}) = \infty$. \square

Our main result for regular problems is given by the following theorem.

THEOREM 5.1. *If*

- (i) $P(s_0, 0) = 0, P'(s_0, 0) \neq 0,$
- (ii) ψ is a normal iteration with order $\rho > 1,$
- (iii) $W^{(0)}(z) = s_0, W^{(1)}(z), \dots, W^{(m)}(z)$ are polynomials in z such that $\text{ord}(E^{(i)}) \geq \rho^i$ for $i = 0, \dots, m,$ where $S(z)$ is the expansion starting with the term $s_0,$
- (iv) $\psi(W^{(i)}(z), W^{(i-1)}(z), \dots, W^{(i-m)}(z)), i = m, m + 1, \dots,$ is a well-defined meromorphic series,

then the iterates

$$W^{(i+1)}(z) \equiv \psi(W^{(i)}(z), W^{(i-1)}(z), \dots, W^{(i-m)}(z)) \pmod{z^{\rho^{i+1}}}$$

satisfy the property that

$$\text{ord}(E^{(i+1)}) \geq \rho^{i+1}, \tag{5.1}$$

and hence the iteration converges.

PROOF. Let $i = m$. By (iv), $\psi W^{(m)}(z), W^{(m-1)}(z), \dots, W^{(0)}(z)$ is a well-defined meromorphic series. Since (4.4) is derived by Taylor series expansions and since the Taylor series expansion is valid over meromorphic series, we have that

$$E^{(m+1)}(z) = \sum_{j_0, \dots, j_m} c_{j_0, \dots, j_m} (E^{(m)}(z))^{j_0} \dots (E^{(0)}(z))^{j_m} \tag{5.2}$$

holds for meromorphic series. The constant term of $P'(S(z), z)$ is $P'(s_0, 0)$, which is nonzero by condition (i). Thus by conditions (ii) and (iii), (5.2) implies that

$$\text{ord}(E^{(m+1)}) \geq \min(j_0 \rho^m + j_1 \rho^{m-1} + \dots + j_m),$$

where the minimum is taken over all the (j_0, \dots, j_m) such that c_{j_0, \dots, j_m} is nonzero for some $P(W, z)$. By the definition of ρ in Section 4, we have

$$\text{ord}(E^{(m+1)}) \geq \rho^{m+1}.$$

By induction, (5.1) can be established for $i = m + 1, m + 2, \dots,$ using similar arguments. The convergence of the iteration follows immediately from Definition 5.2. \square

Remark 5.1. Thus well-defined normal iterations on regular problems always converge. This behavior is strikingly different from the behavior of these iterations on numerical polynomials, where only local convergence is assured unless strong conditions are imposed. Note that the expansion $S(z)$ may converge in only a small disk around the origin; we shall not pursue the domain of convergence here.

Remark 5.2 (5.1) shows that $W^{(i)}$ is a power series with nonnegative powers only rather than a meromorphic series. Until this fact was established it was necessary to work over the field of meromorphic series.

Remark 5.3. Observe that we do not define order for power series valued iteration but only for normal numerical iterations.

Remark 5.4 Note that in Theorem 5.1 we need not assume that $A_n(0) \neq 0$. This fact will be used in the proof of Theorem 6.1.

We apply Theorem 5.1 to two specific iterations. We begin with a Newton-like iteration, which is defined by (5.3) below. This iteration is obtained from the numerical Newton iteration. In the power series setting we hesitate to call it Newton iteration, since Newton [15] actually used a different method for computing the expansion. His

method computes one coefficient per iteration and in general is not as efficient as the Newton-like iteration defined below. We will discuss the Newton-like iteration in some detail since we anticipate it will be the most commonly used iteration in practice. Here and elsewhere we use the notation

$$P'(W, z) \equiv (\partial P / \partial W)(W, z).$$

Recall that the numerical Newton iteration is a normal iteration of order 2. From Theorem 5.1 we have

COROLLARY 5.1.¹ *If*

- (i) $P(s_0, 0) = 0$ and $P'(s_0, 0) \neq 0$,
- (ii) $W^{(0)} = s_0$, i.e. $\text{ord}(E^{(0)}) \geq 1$,

then the iterates $W^{(i)}$ generated by the Newton-like iteration

$$W^{(i+1)}(z) \equiv W^{(i)}(z) - P(W^{(i)}(z), z) / P'(W^{(i)}(z), z) \pmod{z^{2^{i+1}}} \tag{5.3}$$

are well defined and satisfy

$$\text{ord}(E^{(i)}) \geq 2^i \tag{5.4}$$

for $i = 0, 1, 2, \dots$, and hence the iteration converges.

PROOF. We need only show that the iterations $W^{(i)}(z)$ are all well defined. This holds since for all i the constant term in $P'W^{(i)}, z$ is $P'(s_0, 0)$, which is nonzero. \square

Remark 5.5. If we define the valuation of a power series $A(z)$ to be $b^{-\text{ord}(A)}$, where b is any positive constant, then Corollary 5.1 follows from a known theorem in valuation theory (see [2, Ch. II, Theorem 4.2]).

It is easy to show that if $S(z)$ is a polynomial of degree q , then iteration (5.3) will compute it in $\lceil \log_2 q \rceil + 1$ iterations. By a slight modification of the hypotheses of Corollary 5.1 we can replace the inequality (5.4) by equality.

COROLLARY 5.2. *If*

- (i) $P(s_0, 0) = 0, P'(s_0, 0) \neq 0, P''(s_0, 0) \neq 0$,
- (ii) $W^{(0)} = s_0, \text{ord}(E^{(0)}) = 1$,

then the iterates generated by the Newton-like iteration satisfy $\text{ord}(E^{(i)}) = 2^i$. \square

Corollaries 5.1 and 5.2 can easily be generalized to any one-point inverse interpolatory iteration ψ_ρ .

As our second example we consider a secant-like iteration. One has to be somewhat careful in defining this iteration. A straightforward approach would generate iterates by

$$W^{(i+1)} \equiv W^{(i)} - ((W^{(i)} - W^{(i-1)}) / (P(W^{(i)}) - P(W^{(i-1)}))) \cdot P(W^{(i)}) \pmod{z^{\phi^{i+1}}}, \tag{5.5}$$

where $\phi = (1 + \sqrt{5})/2$. Then $W^{(i+1)}$ becomes undefined when $W^{(i)} = W^{(i-1)}$. This happens when there is a "large" gap between the degrees of two consecutive terms, in the expansion which we want to compute. A solution to the problem is given in the following corollary. The idea is to use a perturbed $W^{(i)}$ in (5.5) so that the denominator is guaranteed to be nonzero.

COROLLARY 5.3. *If*

- (i) $P(s_0, 0) = 0, P'(s_0, 0) \neq 0$,
- (ii) $W^{(0)} = s_0, W^{(1)} = s_0 + s_1 z$,

then the iterates $W^{(i)}$ generated by

$$W^{(i+1)} \equiv \bar{W}^{(i)} - ((\bar{W}^{(i)} - W^{(i-1)}) / (P(\bar{W}^{(i)}) - P(W^{(i-1)}))) \cdot P(\bar{W}^{(i)}) \pmod{z^{F_{i+3}}} \tag{5.6}$$

are well defined and satisfy

¹ A result similar to Corollary 5.1 has been proved independently by Professor Lipson [13]

$$\text{ord}(E^{(i)}) \geq F_{i+2},$$

where the F_i is the i -th Fibonacci number (i.e. $F_0 = 0, F_1 = 1$, and $F_{i+1} = F_i + F_{i-1}$) and $\bar{W}^{(i)} = W^{(i)} + z^{F_{i+2}}$.

PROOF. Consider the case $i = 1$. Clearly, $\bar{W}^{(1)} = W^{(1)} + z^2 \neq W^{(0)}$ and $\text{ord}(\bar{W}^{(1)} - W^{(0)}) \leq F_3$. Since by the Taylor series expansion,

$$P(\bar{W}^{(1)}) - P(W^{(0)}) = P'(W^{(0)}) \cdot (\bar{W}^{(1)} - W^{(0)}) + \dots,$$

and since $P'(W^{(0)})$ has a nonzero constant term $P'(s_0, 0)$, we have

$$\text{ord}(P(\bar{W}^{(1)}) - P(W^{(0)})) = \text{ord}(\bar{W}^{(1)} - W^{(0)}) \leq F_3.$$

Hence $(P(\bar{W}^{(1)}) \neq P(W^{(0)}))$. This ensures that $W^{(2)}$ is well defined by (5.6). Note that for $i = 1$, (4.2) holds with $E^{(1)}$ replaced by $\bar{E}^{(1)} = \bar{W}^{(1)} - S$. Thus,

$$\begin{aligned} \text{ord}(P(\bar{W}^{(1)}) - P(\bar{W}^{(0)})) &= \text{ord}(\bar{W}^{(1)} - W^{(0)}) \leq F_3. \\ &\geq \min(\text{ord}(E^{(1)}), F_3) + \text{ord}(E^{(0)}) \geq F_3 + F_2 = F_4. \end{aligned}$$

By induction, one can similarly prove that for $i = 2, 3, \dots$, $W^{(i)}$ is well defined and $\text{ord}(E^{(i)}) \geq F_{i+2}$ \square

Results similar to Corollary 5.3 hold for other suitably modified iterations with memory (i.e. iterations with $m > 0$ in (4.3))

So far we have only dealt with iterations of order greater than one. We now consider an iteration with order one. Define

$$w^{(i+1)} = w^{(i)} - p(w^{(i)})/p'(w^{(i)})$$

for $i = 0, 1, 2, \dots$. Then

$$\begin{aligned} e^{(i+1)} &= e^{(i)} - \frac{p'(s)e^{(i)} + 1/2p''(s)e^{(i)2} + \dots}{p'(s) + p''(s)e^{(i)} + \dots} \\ &= (p''(s)/p'(s))e^{(i)}e^{(i)} - (p''(s)/2p'(s))(e^{(i)})^2 + \sum_{\substack{j+l \geq 3 \\ j \geq 0, l \geq 1}} c_{j,l}(e^{(i)})^j(e^{(i)})^l, \end{aligned} \tag{5.7}$$

where the $c_{j,l}$ are rational expressions whose denominators are powers of $p'(s)$. This implies that the iteration is normal and has order $\rho = 1$. We may use the iteration on power series and obtain the following theorem, which is an easy consequence of (5.7):

THEOREM 5.2 If

- (i) $P(s_0, 0) = 0, P'(s_0, 0) \neq 0,$
- (ii) $W^{(0)} = s_0,$

then the iterates $W^{(i)}$ generated by

$$W^{(i+1)}(z) \equiv W^{(i)}(z) - P(W^{(i)}(z))/P'(W^{(i)}(z)) \pmod{z^{i+2}} \tag{5.8}$$

are well defined and satisfy

$$\text{ord}(E^{(i)}) \geq i + 1,$$

and hence the iteration converges. \square

The iteration (5.8) can be used, for example, to find the initial iterates of an iteration with memory

6. The General Problem: Newton Polygon Process

Recall that our general problem is to compute the value of d and the coefficients s_0, s_1, \dots of an expansion

$$S(z) = \sum_{l=0}^{\infty} s_l \cdot z^{ld}$$

of the algebraic function corresponding to a given

$$P(W, z) = A_n(z)W^n + \dots + A_0(z).$$

In this section we show that the general problem can be reduced to a regular problem by transforming $P(W, z)$ to some $\tilde{P}(W, z)$. The regular problem can then be solved by normal iterations, as described in Section 5

Since $P(s_0, 0) = 0$, s_0 can be obtained by finding a zero of the numerical polynomial $P(W, 0)$. In this section we assume that finding a zero of a numerical polynomial is a primitive operation ² (This assumption will be removed in Section 7 by carrying the zeros symbolically) If $P'(s_0, 0) \neq 0$, we have a regular problem solvable by a normal iteration Hence we assume that $P'(s_0, 0) = 0$. Then s_0 is a multiple zero of the numerical polynomial $P(W, 0)$ and there is more than one expansion of the algebraic function starting with s_0 . We would not expect an iteration starting with $W^{(0)} = s_0$ to converge since the iteration would not "know" to which expansion it should converge. Intuitively the convergence of an iteration requires that it start with an initial segment of a unique expansion. This suggests that we find an initial segment of a unique expansion starting with s_0 . The existence of the segment is guaranteed only if no two expansions coincide, i.e. the discriminant $D(z)$ of $P(W, z)$ with respect to W is not identically equal to zero. Therefore, *in this section we shall assume that $D(z) \neq 0$* . The assumption holds when $P(W, z)$ is irreducible or simply when $P(W, z)$ is square-free [20, Theorem 3.5]. Hence we can make this condition hold by using factorization or square-free decomposition algorithms, but we do not pursue this here.

A classical method for finding an initial segment of a unique expansion uses a geometric aid known as the Newton Polygon, which provides a convenient tool for analyzing a set of inequalities. (Some authors refer to Puiseux's Theorem because of the work of Puiseux [16], but clearly the idea originated with Newton [15, p. 50].) The method has not been subject to algorithmic analysis

We state the Newton Polygon Process adapting, with some modifications, the description in [20] In Theorem 6.1 we show that the Newton Polygon Process transforms the general problem to a regular problem. Theorem 6.1 also gives the connection between the number of identical terms in at least two expansions and the number of Newton Polygon stages Theorem 6.2 gives an a priori bound on the number of stages which differs by at most a factor of two from the optimal bound Example 6.1 shows that in general $P(W, z)$ must be transformed to a new polynomial $\tilde{P}(W, z)$; it is not enough to compute an initial segment of a unique expansion and use it as the initial iterate for a normal iteration on the original polynomial $P(W, z)$.

In the following algorithm let $A_{i,k}(z)$ be the coefficient of W^i in $P_k(W, z)$. If $A_{i,k}(z) \neq 0$, let $a_{i,k}z^{\alpha_{i,k}}$ be the lowest degree term in $A_{i,k}(z)$.

NEWTON POLYGON PROCESS

- N1 $k \leftarrow 1, P_k(W, z) \leftarrow P(W, z)$.
- N2 Plot the points $f_{i,k} = (i, \alpha_{i,k})$ on the xy plane for i such that $A_{i,k}(z) \neq 0$ Join $f_{0,k}$ to $f_{n,k}$ with a convex polygon arc each of whose vertices is an $f_{i,k}$ and such that no $f_{i,k}$ lies below any line extending an arc segment
- N3 If $k = 1$, choose any segment $y + \gamma_k x = \beta_k$ of the arc If $k > 1$, choose a segment with $\gamma_k > 0$. (Such a segment always exists) Let g_k denote the set of indices i for which $f_{i,k}$ lies on the chosen segment Solve the polynomial equation

$$\sum_{i \in g_k} a_{i,k} x^i = 0 \tag{6.1}$$

Let c_k be any of the nonzero roots (Such a nonzero solution always exists)

- N4 If c_k is a simple zero, go to N6, else go to N5
- N5 $P_{k+1}(W, z) \leftarrow z^{-\beta_k} \cdot P_k(z^{\gamma_k} (W + c_k), z), k \leftarrow k + 1$ Go to N2
- N6 $t \leftarrow k$ (Hence t represents the number of stages taken by the Newton Polygon Process)

$$\hat{P}(W, z) \leftarrow z^{-\beta_t} \cdot P_t(z^{\gamma_t} W, z), \tilde{P}(W, z) \leftarrow \hat{P}(W, z^d),$$

² I.e. zeros of a polynomial can be computed to any prespecified precision

where d is the smallest common denominator of $\gamma_1, \dots, \gamma_t$. (γ_1 may be zero. If $\gamma_1 = 0$ we assume that γ_1 has one as its denominator.) Terminate the process

LEMMA 6.1. *After the Newton Polygon Process terminates, the following properties hold:*

- (i) *The coefficients of $\tilde{P}(W, z)$ are polynomials in z .*
- (ii) *c_t is a simple zero of the numerical polynomial $\tilde{P}(W, 0)$.*

PROOF. It is easy to verify (i). To prove (ii) we show that

$$\tilde{P}(W, 0) = \sum_{i \in g_t} a_{i,t} W^i$$

For notational convenience, let $\alpha_{i,t} \equiv \alpha_i$, $a_{i,t} \equiv a_i$, $\beta_t \equiv \beta$, $\gamma_t \equiv \gamma$, $g_t \equiv g$, and let \bar{g} denote the set complementary to g with respect to $\{0, 1, \dots, n\}$. Let

$$P_t(W, z) = (a_n z^{\alpha_n} + Q_n(z)) W^n + \dots + (a_0 z^{\alpha_0} + Q_0(z)),$$

where $\text{ord}(Q_i) > \alpha_i$. Then

$$\hat{P}(W, z) = \sum_{i \in g} a_i W^i + \sum_{j \in \bar{g}} a_j z^{\alpha_j + j\gamma - \beta} W^j + \sum_{i=0}^n z^{i\gamma - \beta} Q_i(z) W^i.$$

Since $\beta = \alpha_i + i\gamma < \alpha_j + j\gamma$, $\forall i \in g$, $\forall j \in \bar{g}$,

$$\tilde{P}(W, 0) = \hat{P}(W, 0) = \sum_{i \in g} a_i W^i. \quad \square$$

THEOREM 6.1. *After the Newton Polygon Process terminates, the following properties hold:*

(i) *The general problem of computing an expansion $S(z)$ of the algebraic function corresponding to $P(W, z)$ has been reduced to the following regular problem: Compute the expansion $\tilde{S}(z)$ starting from c_t for the algebraic function corresponding to $\tilde{P}(W, z)$. Then let*

$$S(z) = \sum_{i=1}^{t-1} c_i z^{\gamma_1 + \dots + \gamma_i} + z^{\gamma_1 + \dots + \gamma_t} \tilde{S}(z^{1/d}).$$

(ii) *$S(z)$ is the unique expansion with starting segment $\sum_{i=1}^t c_i z^{\gamma_1 + \dots + \gamma_i}$.*

(iii) *There is more than one expansion which starts with $\sum_{i=1}^j c_i z^{\gamma_1 + \dots + \gamma_i}$ for every $j < t$. That is, there are at least two expansions which coincide in their first $t - 1$ terms.*

PROOF. By Lemma 6.1, we conclude that the problem of computing $\tilde{S}(z)$ is regular. (Note that the leading coefficient of $\tilde{P}(W, z)$ may vanish at $z = 0$. See Remark 5.4.) (i) follows from $P(W, z) = P(W, z^d)$ and

$$\hat{P}(W, z) = z^{-(\beta_1 + \dots + \beta_t)} P\left(\sum_{i=1}^{t-1} c_i z^{\gamma_1 + \dots + \gamma_i} + z^{\gamma_1 + \dots + \gamma_t} W, z\right).$$

(ii) and (iii) hold since the Newton Polygon Process does not terminate until c_t is a simple zero. \square

Since there is only one expansion which starts with $\sum_{i=1}^t c_i z^{\gamma_1 + \dots + \gamma_i}$, we might expect that if this segment is taken as the initial iterate for a normal iteration then the iteration on the original polynomial $P(W, z)$ rather than on the transformed polynomial $\tilde{P}(W, z)$ will converge. The following example shows this not to be the case; in general we must use the transformed problem

Example 6.1. This problem appears in [9, p. 29], although it is not used to illustrate the point we wish to make here. Let

$$P(W, z) = W^2 - (2 + z + z^3)W + 1 + z + \frac{1}{4}z^2 + z^4.$$

The two expansions are

$$S_1(z) = 1 + 1/2z + z^{3/2} + \dots, \quad S_2(z) = 1 + 1/2z - z^{3/2} + \dots.$$

Suppose that we want to compute $S_1(z)$ by the Newton-like iteration. If we take $W^{(0)} = 1 + 1/2z + z^{3/2}$ in

$$W^{(t+1)} = W^{(t)} - P(W^{(t)}, z)/P'(W^{(t)}, z),$$

we find $W^{(1)} = 1 + 1/2z - 1/4z^{5/2} + \dots$. $W^{(1)}$ differs from S_1 even in the coefficient of $z^{3/2}$! Although there is only one expansion starting with $W^{(0)}$, namely, S_1 , the Newton-like iteration starting from $W^{(0)}$ does not converge to S_1 . \square

We illustrate the Newton Polygon transformation, the transformations of Section 2, and the iterative process with another problem in [9, p. 31].

Example 6.2. Find all the expansions of the algebraic function corresponding to $P(W, z) = -W^3 + zW + z^2$ around $z_0 = \infty$. The first transformation of Section 2 converts $P(W, z)$ to $-z^2W^3 + zW + 1$, which is then converted by another transformation to $-W^3 + zW + z$. The Newton Polygon Process yields $t = 1$, $\beta_1 = 1$, $\gamma_1 = 1/3$, $c_1 = 1$, $d = 3$, and $\bar{P}(W, z) = -W^3 + zW + 1$. Take $W^{(0)} = 1$. Then the Newton-like iteration (5.3) applied to $\bar{P}(W, z)$ gives

$$W^{(1)} = 1 + z/3, \quad W^{(2)} = 1 + z/3 - z^3/81.$$

Thus

$$S(z) = z^{1/3}\bar{S}(z^{1/3}) = z^{1/3} + z^{2/3}/3 - z^{4/3}/81 + \dots.$$

Let $T(z) = S(z)/z = z^{-2/3} + z^{-1/3}/3 - z^{1/3}/81 + \dots$. Then an expansion of the given problem is

$$T(1/z) = z^{2/3} + 1/3z^{1/3} - 1/81z^{-1/3} + \dots.$$

The other two expansions are

$$\theta z^{2/3} + (\theta^2/3)z^{1/3} - (\theta/81)z^{-1/3} + \dots, \quad \theta^2 z^{2/3} + (\theta/3)z^{1/3} - (\theta^2/81)z^{-1/3} + \dots,$$

where θ is the primitive third root of unity. \square

The following theorem gives an a priori bound on the number t of stages in the Newton Polygon Process which differs by at most a factor of two from the optimal bound.

THEOREM 6.2.

$$t \leq \text{ord}(D) + 1. \tag{6.2}$$

Furthermore for all t there exist problems for which $t = 1/2 \text{ord}(D)$.

PROOF. The theorem is trivial if $t = 1$. We assume that $t \geq 2$. Then by (iii) of Theorem 6.1, there are at least two series expansions S_1 and S_2 which agree in the first $t - 1$ nonzero terms. Write

$$S_1 = \sum_{i=1}^{\infty} s_{1,a_i} z^{a_i/d_1}, \quad S_2 = \sum_{i=1}^{\infty} s_{2,b_i} z^{b_i/d_2},$$

where the $\{a_i\}, \{b_i\}$ are strictly increasing nonnegative integer sequences such that none of the s_{1,a_i}, s_{2,b_i} vanish and $s_{1,a_i} = s_{2,b_i}$, $a_i/d_1 = b_i/d_2$ for $i = 1, \dots, t - 1$. Without loss of generality, assume $d_1 \leq d_2$. Note that the cycle which contains S_1 has the series

$$S_{1,j} = \sum_{i=1}^{\infty} s_{1,a_i} \xi_1^{ja_i} z^{a_i/d_1}, \quad j = 0, \dots, d_1 - 1,$$

and the cycle which contains S_2 has the series

$$S_{2,j} = \sum_{i=1}^{\infty} s_{2,b_i} \xi_2^{jb_i} z^{b_i/d_2}, \quad j = 0, \dots, d_2 - 1,$$

where

$$\xi_1 = e^{2\pi\sqrt{-1}/d_1} \quad \text{and} \quad \xi_2 = e^{2\pi\sqrt{-1}/d_2}.$$

Note that we do not rule out the possibility that S_1 and S_2 are in the same cycle and that therefore the cycles $\{S_{1,j}\}$ and $\{S_{2,j}\}$ are identical. Since

$$\xi_1^{a_i} = e^{(2\pi\sqrt{-1}/d_1)ia_i}, \quad \xi_2^{b_i} = e^{(2\pi\sqrt{-1}/d_2)ib_i}$$

and $a_i/d_1 = b_i/d_2$ for $i = 1, \dots, t - 1$, $S_{1,j}$ and $S_{2,j}$ agree in the first $t - 1$ terms for $j = 0, \dots, d_1 - 1$. Hence,

$$\text{ord}(S_{1,j} - S_{2,j}) \geq (b_{t-1} + 1)/d_2 = a_{t-1}/d_1 + 1/d_2.$$

Let

$$V(z) = \prod_{j=0}^{d_1-1} (S_{1,j}(z) - S_{2,j}(z)).$$

Then

$$\text{ord}(D) \geq \text{ord}(V) \geq d_1(a_{t-1}/d_1 + 1/d_2) \geq a_{t-1} + 1.$$

Since the $\{a_i\}$ is a strictly increasing nonnegative integer sequence, $a_{t-1} \geq t - 2$. Thus, $\text{ord}(D) \geq t - 1$, which establishes (6.2). Let

$$S_1(z) = \sum_{j=0}^t z^j, \quad S_2(z) = S_1(z) - z^t,$$

and

$$P(W, z) = (W - S_1(z))(W - S_2(z)).$$

By Theorem 6.1, the Newton Polygon Process has t stages. $\text{ord}(D) = \text{ord}((S_1 - S_2)^2) = 2t$, which completes the proof \square

Theorem 6.2 gives a computable a priori bound but requires the computation of $\text{ord}(D)$. A very cheap bound is given by

COROLLARY 6.1. $t \leq m(2n - 1) + 1$, where $m = \max_{0 \leq i \leq n}(\text{deg } A_i)$.

PROOF. $D(z)$ is a determinant of order $2n - 1$ whose elements are polynomials of maximal degree m . Hence $D(z)$ is a polynomial of degree at most $m(2n - 1)$. Since $D(z)$ cannot vanish identically, $\text{ord}(D) \leq m(2n - 1)$. \square

7. A Symbolic Mode of Computation

The Newton Polygon Process involves computing roots of polynomial equations (6.1). Instead of actually solving the equations, in this section we carry the roots symbolically through their minimum polynomials. We assume that the underlying field Δ is one where exact arithmetic can be performed, such as a finite field or the field Q of rational numbers. Then the expansions can be computed symbolically with exact arithmetic. The following example, where Δ is taken to be Q , will illustrate the idea.

Example 7.1.

$$P(W, z) = W^3 + (z + z^2)W^2 - 2z^2W - 2z^3.$$

We shall compute an expansion of the algebraic function corresponding to $P(W, z)$, using exact rational arithmetic. The first stage of the Newton Polygon Process yields $\gamma_1 = 1$, $\beta_1 = 3$, and $c_1^3 + c_1^2 - 2c_1 - 2 = 0$. Since $c_1^3 + c_1^2 - 2c_1 - 2 = (c_1^2 - 2)(c_1 + 1)$, $c_1 = \sqrt{2}$, $-\sqrt{2}$, or -1 . Suppose that we are interested in the equation starting with $\sqrt{2}$ or $-\sqrt{2}$. Instead of using an approximation to $\sqrt{2}$ or $-\sqrt{2}$, we carry c_1 symbolically through its minimal polynomial $M_1(x) = x^2 - 2$. That is,

$$c_1^2 - 2 = 0. \tag{7.1}$$

Since the equation has only simple zeros, the Newton Polygon Process terminates with $t = 1$, and

$$\tilde{P}(W, z) = z^{-3}P(zW, z) = W^3 + (1 + z)W^2 - 2W - 2.$$

We use the Newton-like iteration (5.3) to compute $\tilde{S}(z)$ such that $\tilde{P}(\tilde{S}(z), z) \equiv 0$. Let $W^{(0)}(z) = c_1$. Then

$$W^{(1)}(z) \equiv c_1 - (c_1^3 - (1 + z)c_1^2 - 2c_1 - 2)/(3c_1^2 + 2(1 + z)c_1 - 2) \pmod{z^2}.$$

Using (7.1), we obtain

$$W^{(1)}(z) = c_1 - 1/3z.$$

Similarly all coefficients of x^j in $W^{(j)}(z)$ can be represented as linear polynomials in c_1 with rational coefficients. By (ii) of Theorem 6.1, a solution to the given problem is

$$S(z) = z\tilde{S}(z) = c_1z - 1/3z^2 + \dots,$$

which represents both the numerical expansions starting with $\sqrt{2}z$ and $-\sqrt{2}z$. \square

In general, when the Newton Polygon Process is performed, $c_k, k = 1, \dots, t$, can be carried symbolically through its minimum polynomial $M_k(x)$ over $Q(c_1, \dots, c_{k-1})$. Then all the coefficients of the expansion $S(z)$ are in the extension field $Q(c_1, \dots, c_t)$. To simplify the computation, one can compute from $M_k(x)$ the minimum polynomial $M(x)$ for c , where c is a primitive element of the extension field $Q(c_1, \dots, c_t)$, i.e. $Q(c) = Q(c_1, \dots, c_t)$. Then the coefficients of the expansion $S(z)$ can all be represented by polynomials of the form $\sum_{i=0}^{h-1} q_i c^i$, where $h = \deg M$ and $q_i \in Q$. $S(z)$ can be computed entirely with exact arithmetic. Furthermore, $S(z)$ gives a simultaneous representation of h numerical expansions; $S(z)$ can be used to produce h numerical expansions by substituting zeros of $M(x)$ for c in the coefficients of $S(z)$. (This implies that $h \leq n$.)

8. Asymptotic Cost Analysis

In this section we analyze the cost of computing the first N terms (including zero terms) of an expansion for large N . Since the Newton Polygon Process is independent of N , by Theorem 6.1 we can without loss of generality assume the problem is regular. Furthermore, since the asymptotic results will be the same for any normal iteration with order greater than one, we shall assume that the iteration (5.3) is used. Our cost measure is the number of operations used over the field Δ . If we carry zeros symbolically as described in Section 7, then we work over an extension field $\Delta(c)$ rather than Δ . If the minimum polynomial for c is of degree h , then operations in $\Delta(c)$ are more expensive than in Δ by a factor of $O(h)$ or $O(h^2)$. Since h is independent of N , in our analysis we shall not be concerned with whether or not zeros of polynomials are carried symbolically.

Let $M(j)$ be the number of operations needed to multiply two j th-degree polynomials over the field Δ . Assume that $M(j)$ satisfies the following mild condition: There are $\alpha, \beta \in (0, 1)$ such that

$$M(\lceil \alpha j \rceil) \leq \beta M(\lceil j \rceil) \tag{8.1}$$

for all sufficiently large j . Observe that $W^{(j)}(z)$ is a polynomial of degree at most $2^j - 1$, and that the computing $W^{(j+1)}(z)$ by (5.3) takes $O(nM(2^j - 1))$ operations. Hence the total cost of computing N terms in the expansion is $O(n(M(N) + M(\lceil N/2 \rceil) + M(\lceil N/4 \rceil) + \dots))$, which is $O(nM(N))$ by condition (8.1). (See [5, Lemma 1.1].) We summarize the result of this section in the following.

THEOREM 8.1 *The first N terms of an expansion of any algebraic function can be computed in $O(nM(N))$ operations over the field Δ . \square*

9 Examples

We choose as our examples calculation of the Legendre polynomials through their generating function, solution of an equation with transcendental coefficients, and calculation of the expansion of a complete elliptic integral. Although the first two

examples are not covered by the theory of this paper, they are covered by easy extensions of our results. Examples 9.1 and 9.3 are illustrations of the many applications of algebraic function expansions.

We use the Newton-like iteration (5.3) in all three examples with the notation:

$$P_i \equiv P(W^{(i)}(z), z), \quad P'_i \equiv P'(W^{(i)}(z), z) \equiv (\partial P/\partial W^{(i)})(W^{(i)}, z^i), \quad \delta_i = p_i/P'_i.$$

Within each iteration step we exhibit enough terms so that $W^{(i)}(z)$ can be computed to $2^i - 1$ terms.

Example 9.1. Legendre polynomials. The generating function for Legendre polynomials,

$$(1 - 2tz + z^2)^{-1/2} = \sum_{i=0}^{\infty} L_i(t)z^i,$$

satisfies

$$P(W, z, t) = (1 - 2tz + z^2)W^2 - 1.$$

Take $W^{(0)} = 1$. Then

$$\begin{aligned} P_0 &= -2tz, & P'_0 &= 2, & \delta_0 &= -tz, & W^{(1)} &= 1 + tz; \\ P_1 &= (1 - 3t^2)z^2 + (2t - 2t^3)z^3, & P'_1 &= 2(1 - tz), & \delta_1 &= 1/2(1 - 3t^2)z^2 + 1/2(3t - 5t^3)z^3, & W^{(2)} &= 1 + tz + 1/2(3t^2 - 1)z^2 + 1/2(5t^3 - 3t)z^3. \end{aligned}$$

Hence the first four Legendre polynomials are

$$L_0(t) = 1, \quad L_1(t) = t, \quad L_2(t) = 1/2(3t^2 - 1), \quad \text{and} \quad L_3(t) = 1/2(5t^3 - 3t).$$

B. Neta, a student at Carnegie-Mellon University, computed the first 32 Legendre polynomials by this iteration using MACSYMA

This example is for illustration; it may not be the best way to compute Legendre polynomials. \square

Example 9.2.

$$P(W, z) = W^2 + (z + 1)W + \sin z.$$

Note that $\sin z = z - z^3/3! + z^5/5! - z^7/7! + \dots$. Take $W^{(0)} = 0$. Then

$$\begin{aligned} P_0 &= z, & P'_0 &= 1, & \delta_0 &= z, & W^{(1)} &= -z; \\ P_1 &= -z^3/6, & P'_1 &= 1, & \delta_1 &= -z^3/6, & W^{(2)} &= -z + z^3/6. \end{aligned} \quad \square$$

Example 9.3. A complete elliptic integral. Define the integral by

$$f(t) = \int_0^{\pi/2} (1 - t^2 \sin^2 \theta)^{-1/2} d\theta.$$

Let

$$P(W, z) = (1 - z)W^2 - 1, \quad z = t^2 \sin^2 \theta.$$

Take $W^{(0)} = 1$. Then

$$\begin{aligned} P_0 &= -z, & P'_0 &= 2, & \delta_0 &= -z/2, & W^{(1)} &= 1 + z/2; \\ P_1 &= -3/4z^2 - 1/4z^3, & P'_1 &= 2 - z, & \delta_1 &= -3/8z^2 - 5/16z^3, \\ & & & & & & W^{(2)} &= 1 + z/2 + 3/8z^2 + 5/16z^3. \end{aligned}$$

$W^{(2)}$ is an initial segment of the algebraic function $S(z)$ corresponding to $P(W, z)$. Since

$$\begin{aligned} f(t) &= \int_0^{\pi/2} S(t^2 \sin^2 \theta) d\theta, \\ f(t) &= \eta_0 + 1/2 \eta_1 t^2 + 3/8 \eta_2 t^4 + 5/16 \eta_3 t^6 + \dots, \end{aligned}$$

where

$$\eta, = \int_0^{\pi/2} \sin^{2i}\theta d\theta.$$

For this simple example the result can be obtained directly by a binomial expansion, but this cannot of course be done in general. \square

10 Extensions

Our aim in this paper has been to show that algebraic functions form an interesting and useful domain in which to do algorithmic and complexity analysis and to exhibit fast algorithms for computing an expansion of an algebraic function. In this initial paper we have restricted ourselves to the “pure” case of algebraic functions where $P(W, z)$ is a polynomial in W with polynomial coefficients. We list some additional problems which we hope to discuss in the future. For a number of these our results (especially on regular problems) apply with minor modifications; others will require major new results.

1. Let W be a scalar variable but take z to be a vector variable. Results similar to those in Section 5 should hold. We have seen this case in Example 9.1.

2. Let the coefficients of $P, A_i(z)$, be power series (rather than polynomials). Results similar to those in Section 5 should hold. See Example 9.2

3. Let both W and z be vector variables. This is the fully multivariate case, which, except for regular problems, is in general very difficult

4. The domain over which we have worked is not algebraically closed since problems with polynomial coefficients lead to solutions represented by fractional power series. If the coefficients are fractional power series, the domain is algebraically closed (Puiseux’s Theorem; see, e.g. [12]), and this is therefore a natural setting. The Newton-like iteration is still valid on fractional power series for regular problems.

5. The field Δ need not be restricted to the complex number field. It is of particular interest to extend all the results to finite fields.

6. An important computational model is the “fully symbolic” one where the coefficients of the expansion series are expressed as functions of the input coefficients.

7. Perform complexity analysis which includes the cost due to the “growth” of coefficients.

ACKNOWLEDGMENTS. We thank H. Woźniakowski, R. Fateman, B. Neta, and A. Werschulz for their comments on the manuscript.

REFERENCES

- 1 AHLFORS, L V *Complex Analysis* McGraw-Hill, New York, sec ed , 1966
- 2 BACHMAN, G *Introduction to P-adic Numbers and Valuation Theory* Academic Press, New York, 1964
- 3 BLISS, G A *Algebraic Functions, Vol XVI* Amer Math Soc Colloquium Publications, Providence, R I , 1933
- 4 BRENT, R P Multiple-precision zero-finding methods and the complexity of elementary function evaluation. In *Analytic Computational Complexity*, J F Traub, Ed , Academic Press, New York, 1976, pp 151-176
- 5 BRENT, R , AND KUNG, H T Fast algorithms for manipulating formal power series. Tech Rep , Comptr Sci Dept , Carnegie-Mellon U , Pittsburgh, Pa , Jan 1976 To appear in *J. ACM*.
- 6 BRIOT, C , AND BOUQUET, J *Théorie des Fonctions Elliptiques* Mallet-Bachelier, Paris, 1859
- 7 EICHLER, M *Introduction to the Theory of Algebraic Numbers and Functions* Transl by G Striker, Academic Press, New York, 1966
- 8 HENSEL, K. *Theorie der Algebraischen Zahlen* Teubner, Berlin, 1908
- 9 JUNG, H W E *Einführung in die Theorie der Algebraischen Funktionen einer Veränderlichen* Walter de Gruyter, Berlin, 1923
- 10 KUNG, H T On computing reciprocals of power series *Numer Math* 22 (1974), 341-348
- 11 KUNG, H T , AND TRAUB, J F Fast algorithms for algebraic functions (abstract) In *Algorithms and Complexity*, J F Traub, Ed , Academic Press, New York, 1976, p 473
- 12 LEFSCHETZ, S *Algebraic Geometry* Princeton U Press, Princeton, N J , 1953

- 13 LIPSON, J. Newton's method A great algebraic algorithm In Proc 1976 ACM Symp on Symbolic and Algebraic Comput., R.D. Jenks, Ed , pp 260-270
- 14 MOSES, J. MACSYMA—The fifth year. SIGSAM Bulletin (ACM) 31 (Aug 1974), 105-110
15. NEWTON, I. Methods of series and fluxions In *The Mathematical Papers of Isaac Newton, Vol III*, D T Whiteside, Ed , Cambridge U Press, 1969
- 16 PUISEUX, V A Recherches sur les fonctions algébriques *J. Math.* 15 (1850), 365-480
- 17 RITT, J F. *Integration in Finite Terms* Columbia U Press, New York, 1948
- 18 SAKS, S , AND ZYGMUND, A *Analytic Functions* American Elsevier, New York, third ed , 1971
- 19 TRAUB, J F *Iterative Methods for the Solution of Equations* Prentice-Hall, Englewood Cliffs, N J , 1964
- 20 WALKER, R J *Algebraic Curves* Princeton U Press, Princeton U , Princeton, N J , 1950
- 21 YUN, D.Y Y Hensel Meets Newton—Algebraic constructions in an analytic setting In *Analytic Computational Complexity*, J F Traub, Ed , Academic Press, New York, 1976, pp 205-216

RECEIVED SEPTEMBER 1976, REVISED JUNE 1977